## Author Index


Achtert, G., see Gielsdorf, W. 351, 357
Ackerman, R., see Kaiser, G. 123
Adachi, K.
— and Asakura, T.
Separation of asymmetrical hybrid hemoglobins by hydrophobic interaction chromatography 303
Adam, M., see Štol, M. 308
Advenier, C., see Naline, E. 177
Ageta, T.
— , Mizobuchi, N. and Kodama, H.
Isotachophoretic analysis for the determination of urinary γ-carboxyglutamic acid 334
Albin, H., see Demotes-Mainard, F. 388
Amacher, D.E.
— and Smith, D.J.
Differences in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 61
Ames, M.M., see Frank, S.K. 225
Andrejak, M., see Hary, L. 396
Andrews, P.A.
— , Bachur, N.R., Musser, S.M., Wright, J. and Callery, P.S.
Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 271
Arnemann, W., see Gielsdorf, W. 351, 357
Asakura, T., see Adachi, K. 303
Assan, R., see Ducrocq, R. 75
Awni, W.M.
— , Clarkson, J. and Guay, D.R.P.
Determination of ciprofloxacin and its 7-ethylenediamine metabolite in human serum and urine by high-performance liquid chromatography 414
Bachur, N.R., see Andrews, P.A. 271
Barlow, R.D.
— , Thompson, P.A. and Stone, R.B.
Simultaneous determination of nicotine, cotinine and five additional nicotine metabolites in the urine of smokers using pre-column derivatisation and high-performance liquid chromatography 375
Bartosevich, J.F., see Kapetanovic, I.M. 458
Bekersky, I., see Puglisi, C.V. 253
Berenson, G.S., see Lee, Y.J. 275
Bergeron, M.G., see Croteau, D. 205
Bhargava, V.O.
Quantitation of 1-(β-D-glucopyranosyl)phenobarbital in plasma and bile 421
Bhavnagri, V.P., see Osman, M.A. 452
Blouin, R.A., see Huang Chandler, M.H. 426
Bonniec, B. Le, see Ducrocq, R. 75
Brassinne, C.
— , Laduron, C., Coune, A., Sculier, J.P., Hollaert, C., Collette, N. and Meunier, F.
High-performance liquid chromatographic determination of amphotericin B in human serum 401
Bredow, J. von, see Kapetanovic, I.M. 458
Brett, M.A.
— , Dierdorf, H.-D., Zussman, B.D. and Coates, P.E.
Determination of paroxetine in human plasma, using high-performance liquid chromatography with fluorescence detection 438
Bron, J., see De Jong, A.P. 233
Callery, P.S., see Andrews, P.A. 271
Carlier, O., see Ducrocq, R. 75
Châtinier, W.M. du, see De Jong, A.P. 233
Chen, S., see Puglisi, C.V. 253
Cheng, L.K., see Osman, M.A. 452
Cingolani, L.
Rapid gas chromatographic determination of phytanic acid from serum of a patient suffering from Refsum's disease 475
Clarkson, J., see Awni, W.M. 414
Coates, P.E., see Brett, M.A. 438
Colistra, C., see Del Principe, D. 329
Collette, N., see Brassinne, C. 401
Corbett, M.E.
— , Scrimgeour, C.M. and Watt, P.W.
Use of *tert.*-butyldimethylsilyl derivatives for gas chromatographic–mass spectrometric analysis of dipeptides 263
Coune, A., see Brassinne, C. 401
Crankshaw, D.P., see Kumar, K. 464
Croteau, D.
— , Vallée, F., Bergeron, M.G. and LeBel, M.
High-performance liquid chromatographic assay of erythromycin and its esters using electrochemical detection 205
Crouzette, J., see Girre, C. 363
Dalferes, Jr., E.R., see Lee, Y.J. 275
Davalli, P., see Desiderio, M.A. 285
DeBethizy, J.D., see Kyerematen, G.A. 191
Decolin, D.
— , Nicolas, A., Motassim, N. and Siest, G.
Direct determination of debrisoquine and 4-hydroxydebrisoquine using a high-perform-

ance liquid chromatographic switching technique 469
De Jong, A.P.
—, Wittebrood, A.J., Du Châtinier, W.M. and Bron, J.
Liquid chromatographic analysis of alzapride and metoclopramide in human plasma and urine using solid-phase extraction 233
De Jong, J.
—, Tjaden, U.R., Visser, E. and Meijer, W.H.
Determination of serotonin and 5-hydroxyindoleacetic acid in urine by reversed-phase ion-pair chromatography with fluorimetric detection 85
Del Principe, D.
—, Colistra, C., Fiorentino, A.M., Spagnolo, G., Mancuso, G. and Menichelli, A.
Detection of fibrinogen heterogeneity by capillary isotachophoresis 329
Demotes-Mainard, F.
—, Vinçon, G., Jarry, C., Necciari, J. and Albin, H.
Micromethod for the determination of cefpiramide in human plasma and urine by high-performance liquid chromatography using automated column switching 388
Desiderio, M.A.
—, Davalli, P. and Perin, A.
Simultaneous determination of γ-aminobutyric acid and polyamines by high-performance liquid chromatography 285
Deyl, Z., see Štol, M. 308
Dierdorf, H.-D., see Brett, M.A. 438
Dieterle, W., see Kaiser, G. 123
DiGiovanni, J.H., see Kapetanovic, I.M. 458
Dubois, J.-P., see Kaiser, G. 123
Du Châtinier, W.M., see De Jong, A.P. 233
Ducrocq, R.
—, Le Bonniec, B., Carlier, O., Assan, R. and Elion, J.
Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 75
Echizen, H., see Takahashi, H. 243
Elion, J., see Ducrocq, R. 75
Emery, P.W., see Woodward, C.J.H. 280
Emm, T.
—, Lesko, L.J. and Perkal, M.B.
Simultaneous determination of doxepin and nordoxepin in serum using high-performance liquid chromatography 445
Er, S.S., see McGarrity, M.J. 37
Fiorentino, A.M., see Del Principe, D. 329
Fishman, R.M., see Moore, G.L. 95
Flatmark, T., see Kvannes, J. 291
Flouvat, B., see Naline, E. 177
Fournier, P.E., see Girre, C. 363
Frank, S.K.
—, Mathiesen, D.A., Whitfield, L.R. and Ames, M.M.
High-performance liquid chromatographic assay for the experimental anticancer agent oxantrazole 225
Friedman, H., see Scavone, J.M. 339
Gielsdorf, W.
—, Arnemann, W., Achtert, G., Hausleiter, H.J. and Li, P.-L.
Determination of timelotem dihydrogenmaleate and its main metabolite N-desmethyltimelotem in plasma by capillary gas chromatography 357
Gielsdorf, W.
—, Hausleiter, H.J., Arnemann, W., Lichtenberger, A. and Achtert, G.
Determination of revenast hydrochloride in human plasma by gas chromatography with nitrogen-selective detection 351
Girre, C.
—, Koch, H., Crouzette, J., Palombo, S. and Fournier, P.E.
Gas chromatographic determination of chlorproethazine in plasma 363
Godeneche, D., see Labarre, P. 381
Gosnell, A.B., see Holder, C.L. 113
Grasela, D.M.
— and Rocci, Jr., M.L.
High-performance liquid chromatographic analysis of pentoxifylline and 1-(5′-hydroxyhexyl)-3,7-dimethylxanthine in whole blood 368
Greenblatt, D.J., see Scavone, J.M. 339
Guay, D.R.P., see Awni, W.M. 414
Guttendorf, R.J., see Huang Chandler, M.H. 426
Hary, L.
— and Andrejak, M.
Dosage du céfotaxime et du désacétylcéfotaxime sériques par chromatographie liquide haute performance avec appariement d'ions 396
Hausleiter, H.J., see Gielsdorf, W. 351, 357
Heiffer, M.H., see Kapetanovic, I.M. 458
Holder, C.L.
—, Korfmacher, W.A., Rushing, L.G., Thompson, Jr., H.C., Slikker, Jr., W. and Gosnell, A.B.
Formation of artifactual metabolites of doxylamine following acid hydrolysis 113

Hollaert, C., see Brassinne, C. 401
Honda, S.
— , Ohmura, T., Segawa, Y., Minematsu, T. and Kakehi, K.
Liquid chromatographic assay of the relative activities of serum pancreatic and salivary α-amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 51
Hsia, J.C., see McGarrity, M.J. 37
Huang Chandler, M.H.
— , Guttendorf, R.J., Blouin, R.A. and Wedlund, P.J.
Simultaneous quantitation of *d*- and *l*-hexobarbital in rat blood by high-performance liquid chromatography 426
Isaksson, K.
— and Kissinger, P.T.
Metabolism of physostigmine in mouse liver microsomal incubations studied by liquid chromatography with dual-electrode amperometric detection 165
Ishizaki, T., see Takahashi, H. 243
Jacobson, K.B., see Yim, J.J. 296
Jarry, C., see Demotes-Mainard, F. 388
Jong, A.P. de, see De Jong, A.P. 233
Jong, J. de, see De Jong, J. 85
Kadofuku, T.
— and Sato, T.
Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 315
Kaiser, G.
— , Ackermann, R., Dieterle, W. and Dubois, J.-P.
Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by gas chromatography–mass spectrometry 123
Kakehi, K., see Honda, S. 51
Kapetanovic, I.M.
— , DiGiovanni, J.H., Bartosevich, J.F., Melendez, V., Schroeder, A.C., Von Bredow, J. and Heiffer, M.H.
Analysis of DL-*threo*-α-(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate (enpiroline), a new candidate antimalarial, in blood using high-performance liquid chromatography 458
Kelly, J.G., see Kinney, C.D. 433
Kim, N., see Yim, J.J. 296
Kinney, C.D.
— and Kelly, J.G.
Liquid chromatographic determination of paracetamol and dextropropoxyphene in plasma 433
Kirch, W., see Santos, S.R.C.J. 155
Kiss, F., see Šalamoun, J. 213
Kissinger, P.T., see Isaksson, K. 165
Koch, H., see Girre, C. 363
Kodama, H., see Ageta, T. 334
Kohout, J., see Kučerová, Z. 323
Korbová, L., see Kučerová, Z. 323
Korfmacher, W.A., see Holder, C.L. 113
Kučerová, Z.
— , Korbová, L. and Kohout, J.
Isolation and characterization of a protein detected in inflamed human gastric mucosa 323
Kumar, K.
— , Morgan, D.J. and Crankshaw, D.P.
Determination of fentanyl and alfentanil in plasma by high-performance liquid chromatography with ultraviolet detection 464
Kvannes, J.
— and Flatmark, T.
Rapid and sensitive assay of ornithine decarboxylase activity by high-performance liquid chromatography of the *o*-phthalaldehyde derivative of putrescine 291
Kyerematen, G.A.
— , Taylor, L.H., DeBethizy, J.D. and Vesell, E.S.
Radiometric–high-performance liquid chromatographic assay for nicotine and twelve of its metabolites 191
Labarre, P.
— , Godeneche, D., Madelmont, J.C., Triana, K., Mathe, G. and Veyre, A.
Qualitative and quantitative analysis of a new sulphur-containing nitrosourea in blood by high-performance liquid chromatography 381
Laudron, C., see Brassinne, C. 401
LeBel, M., see Croteau, D. 205
Le Bonniec, B., see Ducrocq, R. 75
Lee, Y.J.
— , Radhakrishnamurthy, B., Dalferes, Jr., E.R. and Berenson, G.S.
Fractionation of aorta glycosaminoglycans by high-performance liquid chromatography 275
Lesko, L.J., see Emm, T. 445
Li, P.-L., see Gielsdorf, W. 357
Lichtenberger, A., see Gielsdorf, W. 351

Lim, K.T., see Yim, J.J. 296
McClain, C.J., see Schmidt, J. 1
McGarrity, M.J.
—, Er, S.S. and Hsia, J.C.
Hemoglobin-based blood substitutes: characterization of five pyridoxal 5′-phosphate derivatives of hemoglobin 37
Madelmont, J.C., see Labarre, P. 381
Mancuso, G., see Del Principe, D. 329
Mathe, G., see Labarre, P. 381
Mathiesen, D.A., see Frank, S.K. 225
Meijer, W.H., see De Jong, J. 85
Melendez, V., see Kapetanovic, I.M. 458
Meneilly, G.P., see Scavone, J.M. 339
Menichelli, A., see Del Principe, D. 329
Meunier, F., see Brassinne, C. 401
Minder, C.E., see Minder, E.I. 135
Minder, E.I.
—, Schaubhut, R., Minder, C.E. and Vonderschmitt, D.J.
Identification of drugs in human serum by high-performance liquid chromatography with photodiode array detection and a search algorithm for ultraviolet spectra 135
Minematsu, T., see Honda, S. 51
Mizobuchi, N., see Ageta, T. 334
Moore, G.L.
— and Fishman, R.M.
High-performance liquid chromatographic analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 95
Morgan, D.J., see Kumar, K. 464
Motassim, N., see Decolin, D. 469
Musser, S.M., see Andrews, P.A. 271
Naline, E.
—, Flouvat, B., Advenier, C. and Pays, M.
Determination of theophylline and its metabolites in plasma and urine by reversed-phase liquid chromatography using an amine modifier 177
Nambara, T., see Shimada, K. 17
Necciari, J., see Demotes-Mainard, F. 388
Nicolas, A., see Decolin, D. 469
Oe, T., see Shimada, K. 17
Ogata, H., see Takahashi, H. 243
Ohmura, T., see Honda, S. 51
Ohnhaus, E.E., see Santos, S.R.C.J. 155
Osman, M.A.
—, Bhavnagri, V.P., Pinchbeck, F.M., Cheng, L.K. and Wright, G.J.
Determination of duoperone in plasma by high-performance liquid chromatography 452
Palombo, S., see Girre, C. 363
Pao, J., see Puglisi, C.V. 253
Pays, M., see Naline, E. 177
Perin, A., see Desiderio, M.A. 285
Perkal, M.B., see Emm, T. 445
Philipp, E.
Assay for tyrosine hydroxylase in hypothalamic homogenates using high-performance liquid chromatography with electrochemical detection 27
Pinchbeck, F.M., see Osman, M.A. 452
Principe, D. del, see Del Principe, D. 329
Puglisi, C.V.
—, Chen, S., Stelling-Ferrara, M., Pao, J. and Bekersky, I.
Determination of (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthaline, an antiacne agent, and its phenolic metabolite in plasma by reversed-phase high-performance liquid chromatography 253
Radhakrishnamurthy, B., see Lee, Y.J. 275
Roberts, M.S., see Rumble, R.H. 408
Rocci, Jr., M.L., see Grasela, D.M. 368
Rumble, R.H.
— and Roberts, M.S.
High-performance liquid chromatographic assay of the major components of gentamicin in serum 408
Rushing, L.G., see Holder, C.L. 113
Šalamoun, J.
—, Smrž, M., Kiss, F. and Šalamounová, A.
Column liquid chromatography of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 213
Šalamounová, A., see Šalamoun, J. 213
Salari, H.
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 103
Santos, S.R.C.J.
—, Kirch, W. and Ohnhaus, E.E.
Simultaneous analysis of digitoxin and its clinically relevant metabolites using high-performance liquid chromatography and radioimmunoassay 155
Sato, T., see Kadofuku, T. 315
Scavone, J.M.
—, Meneilly, G.P., Greenblatt, D.J. and Friedman, H.
Gas chromatographic analysis of underivatized tocainide 339
Schaubhut, R., see Minder, E.I. 135

Schmidt, J.
— and McClain, C.J.
Separation strategies for *o*-phthalaldehyde–mercaptoethanol derivatives of amino acids for reversed-phase high-performance liquid chromatography 1
Schroeder, A.C., see Kapetanovic, I.M. 458
Scrimgeour, C.M., see Corbett, M.E. 263
Sculier, J.P., see Brassinne, C. 401
Segawa, Y., see Honda, S. 51
Shimada, K.
— , Oe, T. and Nambara, T.
Sensitive ferrocene reagents for derivatization of thiol compounds in high-performance liquid chromatography with dual-electrode coulometric detection 17
Siest, G., see Decolin, D. 469
Slikker, Jr., W., see Holder, C.L. 113
Smith, D.J., see Amacher, D.E. 61
Smrž, M., see Šalamoun, J. 213
Spagnolo, G., see Del Principe, D. 329
Stelling-Ferrara, M., see Puglisi, C.V. 253
Štol, M.
— , Adam, M. and Deyl, Z.
Biospecific affinity chromatography of plasma fibronectin on cross-linked gelatin microbeads 308
Stone, R.B., see Barlow, R.D. 375
Takahashi, H.
— , Ogata, H., Echizen, H. and Ishizaki, T.
Determination of metoclopramide and its glucuronide and sulphate conjugates in human biological fluids (plasma, urine and bile) by ion-pair high-performance liquid chromatography 243
Taylor, L.H., see Kyerematen, G.A. 191
Thompson, P.A., see Barlow, R.D. 375
Thompson, Jr., H.C., see Holder, C.L. 113
Tjaden, U.R., see De Jong, J. 85
Triana, K., see Labarre, P. 381
Uebis, V.
Determination of a new thromboxane $A_2$ receptor blocker in biological fluids by capillary gas chromatography with electron-capture detection 345
Vallée, F., see Croteau, D. 205
Vesell, E.S., see Kyerematen, G.A. 191
Veyre, A., see Labarre, P. 381
Vinçon, G., see Demotes-Mainard, F. 388
Visser, E., see De Jong, J. 85
Von Bredow, J., see Kapetanovic, I.M. 458
Vonderschmitt, D.J., see Minder, E.I. 135
Watt, P.W., see Corbett, M.E. 263
Wedlund, P.J., see Huang Chandler, M.H. 426
Whitfield, L.R., see Frank, S.K. 225
Wittebrood, A.J., see De Jong, A.P. 233
Woodward, C.J.H.
— and Emery, P.W.
Determination of plasma corticosterone using high-performance liquid chromatography 280
Wright, G.J., see Osman, M.A. 452
Wright, J., see Andrews, P.A. 271
Yim, J.J.
— , Lim, K.T., Kim, N. and Jacobson, K.B.
Semi-micro quantitative assay for tryptophan oxygenase by high-performance liquid chromatography 296
Zussman, B.D., see Brett, M.A. 438

## Subject Index

N-Acetyl-L-cysteine
Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 17
Adenine
HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 95
Albumin
Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 75
Alfentanil
Determination of fentanyl and alfentanil in plasma by HPLC with ultraviolet detection 464
Alizapride
LC analysis of alizapride and metoclopramide in human plasma and urine using solid-phase extraction 233
Amines
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285
Amino acids
Separation strategies for o-phthalaldehyde–mercaptoethanol derivatives of amino acids for reversed-phase HPLC 1
γ-Aminobutyric acid
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285
Amphotericin B
HPLC determination of amphotericin B in human serum 401
α-Amylase
LC assay of the relative activities of serum pancreatic and salivary α-amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 51
Angiotensin converting enzyme inhibitor
Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC–MS 123
Antidepressants
Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for ultraviolet spectra 135
Ascorbate-2-phosphate
HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 95
Benzodiazepines
Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for ultraviolet spectra 135
Cadaverine
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285
γ-Carboxyglutamic acid
Isotachophoretic analysis for the determination of urinary γ-carboxyglutamic acid 334
Cefotaxime
Determination of serum cefotaxime and desacetylcefotaxime by ion-pair HPLC 396
Cefpiramide
Micromethod for the determination of cefpiramide in human plasma and urine by HPLC using automated column switching 388
N′-(2-Chloroethyl)-N-[2-(methylsulphinyl)-ethyl]-N′-nitrosourea
Qualitative and quantitative analysis of a new sulphur-containing nitrosourea in blood by HPLC 381
N′-(2-Chloroethyl)-N-[2-(methylsulphonyl)-ethyl]-N′-nitrosourea
Qualitative and quantitative analysis of a new sulphur-containing nitrosourea in blood by HPLC 381
4-[2-(4-Chlorophenylsulphonylamino)ethyl]-phenylacetic acid
Determination of a new thromboxane $A_2$ receptor blocker in biological fluids by capillary GC with electron-capture detection 345
Chlorproethazine
GC determination of chlorproethazine in plasma 363
Chlorpromazine
GC determination of chlorproethazine in plasma 363
Chondroitin sulphate
Fractionation of aorta glycosaminoglycans by HPLC 275
Ciprofloxacin
Determination of ciprofloxacin and its 7-

ethylenediamine metabolite in human serum and urine by HPLC 414
Corticosteroids
Determination of plasma corticosterone using HPLC 280
Corticosterone
Determination of plasma corticosterone using HPLC 280
Cotinine
Simultaneous determination of nicotine, cotinine and five additional nicotine metabolites in the urine of smokers using pre-column derivatisation and HPLC 375
Cytochrome P-450
Difference in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 61
Debrisoquine
Direct determination of debrisoquine and 4-hydroxydebrisoquine using an HPLC switching technique 469
Dermatan sulphate
Fractionation of aorta glycosaminoglycans by HPLC 275
Desacetylcefotaxime
Determination of serum cefotaxime and desacetylcefotaxime by ion-pair HPLC 396
N-Desmethyltimelotem
Determination of timelotem dihydrogenmaleate and its main metabolite N-desmethyltimelotem in plasma by capillary GC 357
Dextropropoxyphene
LC determination of paracetamol and dextropropoxyphene in plasma 433
2,4-Diamino-$N^{10}$-methylpteroid acid
Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 213
Digitoxigenin
Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 155
Digitoxigenin bisdigitoxoside
Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 155
Digitoxigenin monodigitoxoside
Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 155
Digitoxin
Simultaneous analysis of digitoxin and its clinically relevant metabolites using HPLC and radioimmunoassay 155
L-3,4-Dihydroxyphenylalanine
Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 27
1,7-Dimethylxanthine
Determination of theophylline and its metabolites in plasma and urine by reversed-phase HPLC using an amine modifier 177
Dipeptides
Use of *tert.*-butyldimethylsilyl derivatives for GC–MS analysis of dipeptides 263
Doxepin
Simultaneous determination of doxepin and nordoxepin in serum using HPLC 445
Doxylamine
Formation of artifactual metabolites of doxylamine following acid hydrolysis 113
Duoperone
Determination of duoperone in plasma by HPLC 452
Enpiroline
Analysis of DL-*threo*-$\alpha$-(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate (enpiroline), a new candidate antimalarial, in blood using HPLC 458
Enzymes
Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 27
Enzymes
Differences in the constitutive forms of hepatic cytochrome P-450 in male and female adult beagle dogs 61
Enzymes
LC assay of the relative activities of serum pancreatic and salivary $\alpha$-amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 51
Enzymes
Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 291
Enzymes
Semi-micro quantitative assay for tryptophan oxygenase by HPLC 296
Erythromycin
HPLC assay of erythromycin and its esters using electrochemical detection 205
Eseroline
Metabolism of physostigmine in mouse liver

microsomal incubations studied by LC with dual-electrode amperometric detection 165
3-{[1-Ethyoxycarbonyl-3-phenyl-(1*S*)-propyl]-amino}-2,3,4,5-tetrahydro-2-oxo-1-(3*S*)-benzazepine-1-acetic acid
Determination of a new angiotensin converting enzyme inhibitor and its active metabolite in plasma and urine by GC–MS 123
Fentanyl
Determination of fentanyl and alfentanil in plasma by HPLC with ultraviolet detection 464
Ferrocene reagents
Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 17
Fibrinogen
Detection of fibrinogen heterogeneity by capillary isotachophoresis 329
Fibronectin
Biospecific affinity chromatography of plasma fibronectin on cross-linked gelatin microbeads 308
N′-Formylkynurenine
Semi-micro quantitative assay for tryptophan oxygenase by HPLC 296
Gentamicin
HPLC assay of the major components of gentamicin in serum 408
1-(β-D-Glucopyranosyl)phenobarbital
Quantitation of 1-(β-D-glucopyranosyl)phenobarbital in plasma and bile 421
Glucose
Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 75
Glutathione
Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 17
Glycated albumin
Measurement of glycated albumin in diabetic patients by biospecific affinity chromatography 75
Glycoproteins
Biospecific affinity chromatography of plasma fibronectin on cross-linked gelatin microbeads 308
Glycosaminoglycans
Fractionation of aorta glycosaminoglycans by HPLC 275
Haemoglobin
Hemoglobin-based blood substitutes: characterization of five pyridoxal 5′-phosphate derivatives of hemoglobin 37
Haemoglobins
Separation of asymmetrical hybrid hemoglobins by hydrophobic interaction chromatography 303
Heparan sulphate
Fractionation of aorta glycosaminoglycans by HPLC 275
Hexobarbital
Simultaneous quantitation of *d*- and *l*-hexobarbital in rat blood by HPLC 426
Hyaluronic acid
Fractionation of aorta glycosaminoglycans by HPLC 275
4-Hydroxydebrisoquine
Direct determination of debrisoquine and 4-hydroxydebrisoquine using an HPLC switching technique 469
1-(5′-Hydroxyhexyl)-3,7-dimethylxanthine
HPLC analysis of pentoxifylline and 1-(5′-hydroxyhexyl)-3,7-dimethylxanthine in whole blood 368
5-Hydroxyindoleacetic acid
Determination of serotonin and 5-hydroxyindoleacetic acid in urine by reversed-phase ion-pair partition chromatography with fluorimetric detection 85
7-Hydroxymethotrexate
Column LC of methotrexate and its metabolites using a post-column photochemical reactor and fluorescence detection 213
Hypoxanthine
HPLC analysis of ascorbate-2-phosphate, adenine and hypoxanthine in stored human blood 95
Ketones
Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 271
Kynurenine
Semi-micro quantitative assay for tryptophan oxygenase by HPLC 296
Maleimides
Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 17
Maltopentaose, reductively pyridylaminated
LC assay of the relative activities of serum pancreatic and salivary α-amylase using reductively pyridylaminated maltopentaose as a fluorescent substrate 51
Methotrexate
Column LC of methotrexate and its metab-

olites using a post-column photochemical reactor and fluorescence detection 213

3′-Methyl-4-dimethylaminoazobenzene
Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 315

α-Methyldopa
Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 27

Metoclopramide
Determination of metoclopramide and its glucuronide and sulphate conjugates in human biological fluids (plasma, urine and bile) by ion-pair HPLC 243

Metoclopramide
LC analysis of alizapride and metoclopramide in human plasma and urine using solid-phase extraction 233

Neuroleptics
Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for ultraviolet spectra 135

Nicotine
Radiometric–HPLC assay for nicotine and twelve of its metabolites 191

Nicotine
Simultaneous determination of nicotine, cotinine and five additional nicotine metabolites in the urine of smokers using pre-column derivatisation and HPLC 375

Nitrosoureas
Qualitative and quantitative analysis of a new sulphur-containing nitrosourea in blood by HPLC 381

Nordoxepin
Simultaneous determination of doxepin and nordoxepin in serum using HPLC 445

Ornithine decarboxylase
Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 291

Oxantrazole
HPLC assay for the experimental anticancer agent oxantrazole 225

Paracetamol
LC determination of paracetamol and dextropropoxyphene in plasma 433

Paroxetine
Determination of paroxetine in human plasma, using HPLC with fluorescence detection 438

Pentoxifylline
HPLC analysis of pentoxifylline and 1-(5′-hydroxyhexyl)-3,7-dimethylxanthine in whole blood 368

Peptides
Use of *tert.*-butyldimethylsilyl derivatives for GC–MS analysis of dipeptides 263

Phenobarbital
Quantitation of 1-(β-D-glucopyranosyl)phenobarbital in plasma and bile 421

1-Phenyl-1-(2-pyridinyl)ethanol
Formation of artifactual metabolites of doxylamine following acid hydrolysis 113

1-Phenyl-1-(2-pyridinyl)ethylene
Formation of artifactual metabolites of doxylamine following acid hydrolysis 113

Phosphatidylcholine
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 103

Phosphatidylethanolamine
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 103

Phosphatidylinositol
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 103

Phosphatidylserine
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 103

Phospholipids
Comparative study of solid-phase and liquid-phase extraction techniques for isolation of phospholipids from plasma 103

Physostigmine
Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 165

Phytanic acid
Rapid GC determination of phytanic acid from serum of a patient suffering from Refsum's disease 475

Polyamines
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285

Proteins
Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 315

Proteins, gastric mucosa
Isolation and characterization of a protein

detected in inflamed human gastric mucosa 323
Putrescine
Rapid and sensitive assay of ornithine decarboxylase activity by HPLC of the *o*-phthalaldehyde derivative of putrescine 291
Putrescine
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285
Pyridoxal 5′-phosphate-labelled haemoglobin
Hemoglobin-based blood substitutes: characterization of five pyridoxal 5′-phosphate derivatives of hemoglobin 37
Revenast hydrochloride
Determination of revenast hydrochloride in human plasma by GC with nitrogen-selective detection 351
Rubreserine
Metabolism of physostigmine in mouse liver microsomal incubations studied by LC with dual-electrode amperometric detection 165
Serotonin
Determination of serotonin and 5-hydroxyindoleacetic acid in urine by reversed-phase ion-pair partition chromatography with fluorimetric detection 85
Serum proteins
Detection of the changes in serum proteins during chemical hepatocarcinogenesis by two-dimensional electrophoresis under non-denaturing conditions 315
Silylated N,O-ketals
Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 271
Spermidine
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285
Spermine
Simultaneous determination of γ-aminobutyric acid and polyamines by HPLC 285
Steroids
Determination of plasma corticosterone using HPLC 280
(*E*)-1,2,3,4-Tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene
Determination of (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene, an antiacne agent, and its phenolic metabolite in plasma by reversed-phase HPLC 253
(*E*)-4-[2-(5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-2-methylethenyl]-phenol
Determination of (*E*)-1,2,3,4-tetrahydro-1,1,4,4-tetramethyl-6-(1-methyl-2-phenylethenyl)naphthalene, an antiacne agent, and its phenolic metabolite in plasma by reversed-phase HPLC 253
Theophylline
Determination of theophylline and its metabolites in plasma and urine by reversed-phase HPLC using an amine modifier 177
Thiol compounds
Sensitive ferrocene reagents for derivatization of thiol compounds in HPLC with dual-electrode coulometric detection 17
Thromboxane $A_2$ receptor blocker
Determination of a new thromboxane $A_2$ receptor blocker in biological fluids by capillary GC with electron-capture detection 345
Timelotem dihydrogenmaleate
Determination of timelotem dihydrogenmaleate and its main metabolite N-desmethyltimelotem in plasma by capillary GC 357
Tocainide
GC analysis of underivatized tocainide 339
N-Trimethylsilylimidazole
Silylated N,O-ketals from the reaction of ketones with N-trimethylsilylimidazole 271
Tryptophan oxygenase
Semi-micro quantitative assay for tryptophan oxygenase by HPLC 296
Tyrosine hydroxylase
Assay for tyrosine hydroxylase in hypothalamic homogenates using HPLC with electrochemical detection 27
Ultraviolet spectra, search algorithm
Identification of drugs in human serum by HPLC with photodiode array detection and a search algorithm for ultraviolet spectra 135
Uric acids
Determination of theophylline and its metabolites in plasma and urine by reversed-phase HPLC using an amine modifier 177
Xanthine metabolites
Determination of theophylline and its metabolites in plasma and urine by reversed-phase HPLC using an amine modifier 177